

Feedlot 40-20 R400

SUPPLEMENT FOR BEEF CATTLE



Specie: Beef

Type of Feed: Protein supplement

Form of Feed: Pellet

General Description: Designed to improve feed efficiency in growing and finishing beef cattle.

PROTEIN SUPPLEMENT FOR BEEF CATTLE

Guaranteed Analysis

Crude Protein	(not less than)	40.0%
Crude Fat	(not less than)	2.0%
Crude Fiber	(not more than)	12.0%
Calcium	(not less than)	7.0%
Calcium	(not more than)	8.0
Phosphorus	(not more than)	0.50%
Salt	(not less than)	3.5%
Salt	(not more than)	4.5%
Magnesium	(not less than)	0.8%
Copper	(not less than)	190 ppm
Selenium	(not less than)	4.1 ppm
Zinc	(not less than)	650 ppm
Vitamin A	(not less than)	40,000 IU/lb
Vitamin D3	(not less than)	10,000 IU/lb
Vitamin E	(not less than)	50 IU/lb

Ingredients:

Processed Grain By-Products, Calcium Carbonate, Plant Protein Products, Urea, Potassium Chloride, Salt, Magnesium Oxide, Sodium Selenite, Zinc Sulfate, Roughage Products, Manganese Sulfate, Manganous Oxide, Zinc Oxide, Basic Copper Chloride, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Copper Sulfate, Potassium Iodide, Mineral Oil, Cobalt Carbonate, Ethylenediamine Dihyridide

Feeding Directions: Feed to cattle at the rate of 1.0-1.5 lbs per head per day, mix with grain or with roughage. Always provide clean water at all times. CAUTION: Do NOT allow horses or other equine access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Consumption by unapproved species in toxic reactions. Do NOT feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use. DO NOT exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. A withdrawal time has not been established for pre-ruminating calves. Do NOT use in calves to be processed for veal.